

QFLOW-MRI[®] References

Validation Studies

- **Geest RJ van der, Niezen RA, Wall EE van der, Roos A de, Reiber JHC.**
Automated measurement of volume flow in the ascending aorta using MR velocity maps: Evaluation of inter- and interobserver variability in healthy volunteers.
J Comp Assist Tomogr 1998; 22 (6): 904-11.
- **Onen F, Feugeas MCH, Marco G de, Baron G, Ravaud P, Legrain S, Moretti JL, Schouman Claeys E, Peretti II.**
Cerebrospinal fluid MR dynamics and risk of falls in the elderly.
J Neuroradiol 2005; 32: 3-9
- **Salm LP, Schuijf JD, Lamb HJ, Bax JJ, Vliegen HW, Jukema JW, Wall EE van der, Roos A de, Doornbos J.**
Validation of a high-resolution, phase contrast cardiovascular magnetic resonance sequence for evaluation of flow in coronary artery bypass grafts.
J Cardiovasc Magn Reson 2007; 9: 557-563.

Other Publications:

1. **Hoogendoorn LI, Pattynama PMT, Buis B, Geest RJ van der, Wall EE van der, Roos A de.**
Noninvasive evaluation of aortocoronary bypass grafts with magnetic resonance flow mapping.
Am J Cardiol 1995; 75: 845-48.
2. **Helbing WA, Bosch JG, Maliepaard C, et al.**
Comparison of echocardiographic methods with magnetic resonance imaging for assessment of right ventricular function in children.
Am J Cardiol 1995; 76 :589-94.
3. **Hundley WG, Li HF, Willard JE, et al.**
Magnetic resonance imaging assessment of the severity of mitral regurgitation: Comparison with invasive techniques.
Circulation 1995; 92: 1151-58.
4. **Geest RJ van der, Buller VGM, Reiber JHC.**
Automated quantification of flow velocity and volume in the ascending and descending aorta using MR flow velocity mapping.
IEEE, Computers in Cardiology 1995: 29-32.

- 5. Helbing WA, Niezen RA, le Cessie S, Geest RJ van der, Ottenkamp J, Roos A de.**
Right ventricular diastolic function in children with pulmonary regurgitation after repair of Tetralogy of Fallot: Volumetric evaluation by magnetic resonance velocity mapping.
J Am Coll Cardiol 1996; 28:1827-35.
- 6. Niezen RA, Helbing WA, Geest RJ van der, Rebergen SA, Roos A de.**
Biventricular systolic function and mass studied with MR imaging in children with pulmonary regurgitation after repair for Tetralogy of Fallot.
Radiology 1996; 201: 135-40.
- 7. Kayser HWM, Stoel BC, Wall EE van der, Geest RJ van der, Roos A de.**
MR velocity mapping of tricuspid flow: Correction for through-plane motion.
J Magn Reson Imag 1997; 7 (4): 669-73.
- 8. Kayser HWM, SchaliJ MJ, Wall EE van der, Stoel BC, Roos A de.**
Biventricular function in patients with nonischemic right ventricular tachyarrhythmias assessed with MR imaging.
AJR 1997; 169 (4): 995-9.
- 9. Roos A de, Helbing WA, Niezen RA, Rebergen SA, Wall EE van der, Ottenkamp J.**
Magnetic resonance imaging in adult congenital heart disease.
In: Current and future applications of magnetic resonance in cardiovascular disease. Higgins CB, Inwall JS, Pohost GM, editors. Futura Publishing Company, Inc. Armonk, NY 1998: 163-72.
- 10. Westenberg JJM, Wasser MNJM, Geest RJ van der, et al.**
Variations in blood flow waveforms in stenotic renal arteries by 2D phase-contrast Cine MRI.
J Magn Reson Imag 1998; 8: 590-7.
- 11. Geest RJ van der, Niezen RA, Wall EE van der, Roos A de, Reiber JHC.**
Automated measurement of volume flow in the ascending aorta using MR velocity maps: evaluation of inter- and intraobserver variability in healthy volunteers.
J Comput Assisted Tomography 1998; 22: 904-11.
- 12. Niezen RA, Doornbos J, Wall EE van der, Roos A de.**
Measurement of aortic and pulmonary flow with MRI at rest and during physical exercise.
J Comput Assist Tomogr 1998; 22 (2):194-201.
- 13. Kozerke S, Botnar R, Oyre S, Scheidegger MB, Pedersen EM, Boesiger P.**
Automatic vessel segmentation using active contours in cine phase contrast flow measurements.

J Magn Reson Imag 1999; 10 (1): 41-51.

14. Marcus JT, Smeenk HG, Kuijer JP, Geest RJ van der, Heethaar RM, Rossum AC van.

Flow profiles in the left anterior descending and the right coronary artery assessed by MR velocity quantification: effects of through-plane and in-plane motion of the heart.
J Comput Assist Tomogr 1999; 23 (4): 567-76.

15. Geest RJ van der, Reiber JHC.

Quantification in cardiac MRI.
J Magn Reson Imag 1999;10 (5): 602-8.

16. Langerak SE, Kunz P, Vliegen HW, Lamb HJ, Jukema JW, Wall EE van der, Roos A de.

Improved MR Flow mapping in coronary artery bypass grafts during adenosine-induced stress.
Radiology 2001; 218: 540-7.

17. Groenink, M, Roos A de, Mulder BJM, Verbeeten B, Timmermans J, Zwinderman AH, Spaan JAE, Wall EE van der.

Biophysical properties of the normal-sized aorta in patients with Marfan syndrome: evaluation with MR Flow mapping.
Radiologie 2001; 219: 535-40.

18. Beyerbacht HP, Lamb HJ, Laarse A van der, Vliegen HW, Leujes F, Hazekamp MG, Roos A de, Wall EE van der.

Aortic valve replacement in patients with aortic valve stenosis improves myocardial metabolism and diastolic function.
Radiologie 2001; 219: 637-43.

19. Bedaux WLF, Hofman MBM, Visser CA, Rossum, AC van.

Simultaneous noninvasive measurement of blood flow in the great cardiac vein and left anterior descending artery.
J Magn Reson Imag 2001; 3 (3): 227-35.

20. Langerak SE, Kunz P, Vliegen HW, Jukema JW, Zwinderman AH, Steendijk P, Lamb HJ, Wall EE van der, Roos A de.

MR Flow mapping in coronary artery bypass grafts: a validation study with Doppler flow measurements¹.
Radiology 2002; 222 (1): 127-35.

21. Bedaux WLF, Hofman MBM, Vyt SLA, Bronzwaer JGF, Visser CA, Rossum AC van.

Assessment of coronary artery bypass graft disease using cardiovascular magnetic resonance determination of flow reserve.

J Am Coll Cardiol 2002; 40(10): 1848-55.

- 22. Roest AAW, Roos A de, Lamb HJ, Helbing WA, Aardweg JG van den, Doornbos J, Wall EE van der, Kunz P.**
Tetralogy of Fallot: postoperative delayed recovery of left ventricular stroke volume after physical exercise - Assessment with fast MR imaging.
Radiology 2003; 226(1): 278-84.
- 23. Wong P, Graves MJ, Lomas DJ**
Integrated physiological flow simulator and pulse sequence monitoring system for MRI
Med Biol Eng Comput 2008; 46: 399-406
- 24. Rigsby CK, Hilpiper N, McNeal GR, Zhang G, Boylan EE, Popescu AR, Choi G, Greiser A, Deng J**
Analysis of an automated background correction method for cardiovascular MR phase contrast imaging in children and young adults.
Pediatr Radiol 2013; DOI 10.1007/s00247-013-2830-y
- 25. Kozak MF, Yoo S-J, Mertens L, Ho A, Grosse-Wortmann L**
Assessment of ductal blood flow in newborns with obstructive left heart lesions by cardiovascular magnetic resonance
J Cardiovasc Magn Reson 2013; 15: 45
- 26. Joseph AA**
Real-time MRI of moving spins using undersampled radial FLASH
PhD thesis, University of Würzburg, 2013

- 27. Flacke S, Allen JS, Chia JM, Wible JH, Periasamy MP, Adams MD, Adzamli IK, Lorenz CH.**
Characterization of viable and nonviable myocardium at MR imaging: comparison of gadolinium-based extracellular and blood pool contrast materials versus manganese-based contrast materials in a rat myocardial infarction model.
Radiology 2003; 226(3): 731-8.
- 28. Langerak SE, Vliegen HW, Jukema JW, Kunz P, Zwinderman AH, Lamb HJ, Wall EE van der, Roos A de.**
Value of magnetic resonance imaging for the noninvasive detection of stenosis in coronary artery bypass grafts and recipient coronary arteries.
Circulation 2003; 107(11): 1502-8.
- 29. Langerak SE, Vliegen HW, Jukema JW, Zwinderman AH, Lamb HJ, Roos A de, Wall EE van der.**
Vein graft function improvement after percutaneous intervention: evaluation with MR Flow mapping.
Radiology 2003; 228 (3): 834-41.
- 30. Hout RJ van den, Lamb HJ, Aardweg JG van den, Schot R, Steendijk P, Wall EE van der, Bax JJ, Roos A de.**
Real-time MR imaging of aortic flow: influence of breathing on left ventricular stroke volume in chronic obstructive pulmonary disease.
Radiology 2003; 229 (2): 513-9.
- 31. Paelinck BP, Lamb HJ, Bax JJ, Wall EE van der, Roos A de.**
MR flow mapping of dobutamine-induced changes in diastolic heart function.
J Magn Reson Imag 2004; 19 (2): 176-81.
- 32. Nollen GJ, Groenink M, Tijssen JGP, Wall EE van der, Mulder BJM.**
Aortic stiffness and diameter predict progressive aortic dilatation in patients with Marfan syndrome.
Eur Heart J 2004; 25 (13): 1146-52.
- 33. Salm LP, Langerak SE, Vliegen HW, Jukema JW, Bax JJ, Zwinderman AH, Wall EE van der, Roos A de, Lamb HJ.**
Blood flow in coronary artery bypass vein grafts: volume versus velocity at cardiovascular MR imaging.
Radiology 2004; 232 (3): 915-20.
- 34. Salm LP, Bax JJ, Vliegen HW, Langerak SE, Dibbets P, Jukema JW, Lamb HJ, Pauwels EKJ, Roos A de, Wall EE van der.**
Functional significance of stenoses in coronary artery bypass grafts. Evaluation by single-photon emission computed tomography perfusion imaging, cardiovascular magnetic resonance, and angiography.
JACC 2004; 44 (9): 1877-82.

- 35. Straten A van, Vliegen HW, Hazekamp MG, Bax JJ, Schoof PH, Ottenkamp J, Wall EE van der, Roos A de.**
Right ventricular function after pulmonary valve replacement in patients with tetralogy of fallot.
Radiology 2004; 233(3): 824-9.
- 36. Westenberg JJM, Danilouchkine MG, Doornbos J, Bax JJ, Geest RJ van der, Labadie G, Lamb HJ, Versteegh MIM, Roos A de, Reiber JHC.**
Accurate and reproducible mitral valvular blood flow measurement with three-directional velocity-encoded magnetic resonance imaging.
J Cardiovasc Magn Reson 2004; 6 (4): 767-76.
- 37. Paelinck BP, Roos A de, Bax JJ, Bosmans JM, Geest RJ van der, Dhondt D, Parizel PM, Vrints CJ, Lamb HJ.**
Feasibility of tissue magnetic resonance imaging. A pilot study in comparison with tissue Doppler imaging and invasive measurement.
JACC 2005; 45(7): 1109-16.
- 38. Dam VH ten, Box FMA, Craen AJM de, Heuvel DMJ van den, Bollen ELEM, Murray HM, Buchem MA van, Westendorp RGJ, Blauw GJ on behalf of the PROSPER Study Group.**
Lack of effect of pravastatin on cerebral blood flow or parenchymal volume loss in elderly at risk for vascular disease.
Stroke 2005; 36: 1633-6.
- 39. Pul C van, Jong NMCM de, Beek LM van, Pasmans HLM, Wijn PFF, Visser RF.**
MRI for diagnosing aortic valve stenosis: a comparison study of MRI and ultrasound.
Neth Heart J 2005; 13 (11): 387-92.
- 40. Straten A van, Vliegen HW, Lamb HJ, Roes SD, Wall EE van der, Hazekamp MG, Roos A de.**
Time course of diastolic and systolic function improvement after pulmonary valve replacement in adult patients with tetralogy of fallot.
JACC 2005; 46(8):1559-64.
- 41. Oshinski JN, Curtin JL, Loth F.**
Mean-average wall shear stress measurements in the common carotid artery.
J Cardiovasc Magn Reson 2006; 8: 717-22.
- 42. Grotenhuis HB, Ottenkamp J, Westenberg JJM, Bax JJ, Kroft LJM, Roos A de.**
Reduced aortic elasticity and dilatation are associated with aortic regurgitation and left ventricular hypertrophy in nonstenotic bicuspid aortic valve patients.
JACC 2007; 49(15): 1660-5.

- 43. Dam VH ten, Heuvel DMJ van den, Craen AJM de, Bollen ELEM, Murray HM, Westendorp RGJ, Blauw GJ, Buchem MA van.**
Decline in total cerebral blood flow is linked with increase in periventricular but not deep white matter hyperintensities.
Radiology 2007;243(1):198-203.
- 44. Salm LP, Schuijf JD, Lamb HJ, Bax JJ, Vliegen HW, Jukema JW, Wall EE van der, Roos A de, Doornbos J.**
Validation of a high-resolution, phase contrast cardiovascular magnetic resonance sequence for evaluation of flow in coronary artery bypass grafts.
J Cardiovasc Magn Reson 2007; 9: 557-563.
- 45. Meer RW van der, Diamant M, Westenberg JJM, Doornbos J, Bax JJ, Roos A de, Lamb HJ.**
Magnetic resonance assessment of aortic pulse wave velocity, aortic distensibility, and cardiac function in uncomplicated type 2 diabetes mellitus.
J Cardiovasc Magn Reson 2007; 9: 645-651.
- 46. Westenberg JJM, Lamb HJ, Geest RJ van der, Bleeker GB, Holman ER, Schalijs MJ, Roos A de, Wall EE van der, Reiber JHC, Bax JJ.**
Assessment of left ventricular dyssynchrony in patients with conduction delay and idiopathic dilated cardiomyopathy. Head-to-head comparison between tissue doppler imaging and velocity-encoded magnetic resonance imaging.
JACC 2006;47(10):2042-8.
- 47. Grotenhuis HB, Kroft LJM, Elderen SGC van, Westenberg JJM, Doornbos J, Hazekamp MG, Vliegen HW, Ottenkamp J, Roos A de.**
Right ventricular hypertrophy and diastolic dysfunction in arterial switch patients without pulmonary artery stenosis.
Heart 2007;93:1604-1608.
- 48. Wolferen SA van, Marcus JT, Westerhof N, Spreeuwenberg MD, Marques KMJ, Bronzwaer JGF, Henkens IR, Gan CTJ, Boonstra A, Postmus PE, Vonk-Noordegraaf A.**
Right coronary artery flow impairment in patients with pulmonary hypertension.
Eur Heart J 2008;29:120-127.
- 49. Grotenhuis HB, Roos A de, Ottenkamp J, Schoof PH, Vliegen HW, Kroft LJM.**
MR imaging of right ventricular function after the Ross procedure for aortic valve replacement: initial experience.
Radiology 2008;246(2):394-400.
- 50. Johnson K, Sharma P, Oshinski J.**
Coronary artery flow measurement using navigator echo gated phase contrast magnetic resonance velocity mapping at 3.0 T.
J Biomechanics 2008;41:595-602.

- 46. Meer RW van der, Hammer S, Smit JWA, Frölich M, Bax JJ, Diamant M, Rijzewijk LJ, Roos A de, Romijn JA, Lamb HJ.**
Short-term caloric restriction induces accumulation of myocardial triglycerides and decreases left ventricular diastolic function in healthy subjects.
Diabetes 2007;56:2849-2853.
- 47. Hammer S, Meer RW van der, Lamb HJ, Schär M, Roos A de, Smit JWA, Romijn JA.**
Progressive caloric restriction induces dose dependent changes in myocardial triglyceride content and diastolic function in healthy men.
J Clin Endocrin Metab 2007;2007-2015.
- 48. Westenberg JJM, Braun J, Veire NR van de, Klautz RJM, Versteegh MIM, Roes SD, Geest RJ van der, Roos A de, Wall EE van der, Reiber JHC, Bax JJ, Dion RAE.**
Magnetic resonance imaging assessment of reverse left ventricular remodeling late after restrictive mitral annuloplasty in early stages of dilated cardiomyopathy.
J Thorac Cardiovasc Surg 2008;135:1247-53.
- 49. Meer RW van der, Rijzewijk LJ, Diamant M, Hammer S, Schär M, Bax JJ, Smit JWA, Romijn JA, Roos A de, Lamb HJ.**
The ageing male heart: myocardial triglyceride content as independent predictor of diastolic function.
Eur Heart J 2008;29:1516-1522.
- 50. Grotenhuis HB, Ottenkamp J, Fontein D, Vliegen HW, Westenberg JJM, Kroft LJM, Roos A de.**
Aortic elasticity and left ventricular function after arterial switch operation: MR imaging – initial experience.
Radiol 2008;249(3):801-809.
- 51. Rijzewijk LJ, Meer RW van der, Smit JWA, Diamant M, Bax JJ, Hammer S, Romijn JA, Roos de A, Lamb HJ.**
Myocardial steatosis is an independent predictor of diastolic dysfunction in type 2 diabetes mellitus.
JACC 2008;52(22):1793-9.
- 52. Westenberg JJM, Roes SD, Marsan NA, Binnendijk NMJ, Doornbos J, Bax JJ, Reiber JHC, Roos A de, Geest RJ van der.**
Mitral valve and tricuspid valve blood flow: accurate quantification with 3D velocity-encoded MR imaging with retrospective valve tracking.
Radiol 2008;249(3):792-800.

- 53. Dockum WG van, Knaapen P, Hofman MBM, Kuijer JPA, Cate FJ ten, Berg JM ten, Beek AM, Twisk JWR, Rossum AC van.**
Impact of alcohol septal ablation on left anterior descending coronary artery blood flow in hypertrophic obstructive cardiomyopathy.
Int J Cardiovasc Imaging 2009;25:511-518.
- 54. Romeih S, Kroft LJ, Bokenkamp R, Schalijs MJ, Grotenhuis H, Hazekamp MG, Groenink M, Roos A de, Blom NA.**
Delayed improvement of right ventricular diastolic function and regression of right ventricular mass after percutaneous pulmonary valve implantation in patients with congenital heart disease.
Am Heart J 2009;158:40-6.
- 55. Grotenhuis HB, Ottenkamp J, Bruijn L de, Westenberg JJM, Vliegen HW, Kroft LJM, Roos A de.**
Aortic elasticity and size are associated with aortic regurgitation and left ventricular dysfunction in tetralogy of Fallot after pulmonary valve replacement.
Heart 2009;95:1931-1936.
- 56. Shanks M, Siebelink HMJ, Delgado V, Veire NRL van de, Ng ACT, Sieders A, Schuijff JD, Lamb HJ, Marsan NA, Westenberg JJM, Kroft LJ, Roos A de, Bax JJ.**
Quantitative assessment of mitral regurgitation. Comparison between three-dimensional transesophageal echocardiography and magnetic resonance imaging.
Circ Cardiovasc Imaging 2010;3:694-700
- 57. Hickson SS, Butlin M, Graves M, Taviani V, Avolio AP, McEniery CM, Wilkinson IB.**
The relationship of age with regional aortic stiffness and diameter.
JACC Cardiovasc Imaging 2010;3:1247-1255
- 58. Hays AG, Hirsch GA, Kelle S, Gerstenblith G, Weiss RG, Stuber M.**
Noninvasive visualization of coronary artery endothelial function in healthy subjects and in patients with coronary artery disease.
JACC 2010;56(20):1657-65
- 59. Taviani V, Patterson AJ, Graves MJ, Hardy CJ, Worters P, Sutcliffe MPF, Gillard JH**
Accuracy and repeatability of Fourier velocity encoded M-mode and two-dimensional cine phase contrast for pulse wave velocity measurement in the descending aorta
J Magn Reson Imaging 2010; 31: 1185-1194.
- 60. Hulst AE van der, Westenberg JJM, Kroft LJM, Bax JJ, Blom NA, Roos A de, Roest AAW**

Tetralogy of Fallot: 3D velocity-encoded MR imaging for evaluation of right ventricular valve flow and distolic function in patients after correction
Radiology 2010; 256: 724-734

- 61. Elderen SGC van, Brandts A, Grond J van der, Westenberg JJM, Kroft LJM, Buchem MA van, Smit JWA, Roos A de.**
Cerebral perfusion and aortic stiffness are independent predictors of white matter brain atrophy in type 1 diabetic patients assessed with magnetic resonance imaging.
Diabetes Care 2011; 34:459-463
- 62. Rüssel IK, Brouwer WP, Germans Tj, Knaapen P, Marcus JT, Velden J van der, Götte MJW, Rossum AC van**
Increased left ventricular torsion in hypertrophic cardiomyopathy mutation carriers with normal wall thickness.
J Cardiovasc Magn Reson 2011; 13:3
- 63. Marsan NA, Westenberg JJM, Roes SD, Bommel RJ van, Delgado V, Geest RJ van der, Roos A de, Klautz RJ, Reiber JC, Bax JJ**
Three-dimensional echocardiography for the preoperative assessment of patients with left ventricular aneurysm
Ann Thorac Surg 2011; 91: 113-122.
- 64. Bonekamp D, Degaonkar M, Barker PB.**
Quantitative cerebral blood flow in dynamic susceptibility contrast MRI using total cerebral flow from phase contrast magnetic resonance angiography.
Magnetic Res. in Medicine (2011); 66:57-66
- 65. Brouwer WP, Germans T, Head MC, Velden J van der, Heymans MW, Christiaans I, Houweling AC, Wilde AA, Rossum AC van**
Multiple myocardial crypts on modified long-axis view are a specific finding in pre-hypertrophic HCM mutation carriers.
Eur Heart J-Cardiovasc Imaging 2012; 13: 292-297
- 66. Al Nafisi B, Amerom JFP van, Forsey J, Jaeggi E, Grosse-Wortmann L, Yoo S-J, Macgowan CK, Seed M**
Fetal circulation in left-sided congenital heart disease measured by cardiovascular magnetic resonance: a case-control study
J Cardiovasc Magn Reson 2013; 15: 65 doi: 10.1186/1532-429X-15-65
- 67. Joseph A, Kowallick JT, Merboldt K-D, Voit D, , Schaetz S, Zhang S, Sohns JM, Lotz J, Frahm J.**
Real-Time Flow MRI of the aorta at a resolution of 40 msec
JMRI 2013
- 68. Turkbey EB, Jain A, Johnson C, Redheuil A, Arai A, Gomes AS, Carr J, Hundley WG, Teixido-Tura G, Eng J, Lima JAC, Bluemke DA**

Determinants and normal values of ascending aortic diameter by age, gender, and race/ethnicity in the Multi-Ethnic Study of Atherosclerosis (MESA)
J Magn Reson Imaging 2014; 39: 360-368.

- 69. Kröner ESJ, Lamb HJ, Siebelink H-MJ, Putter H, Geest RJ van der, Wall EE van der, Roos A de, Westenberg JJM.**
Coupling of vessel wall morphology and function in the aorta and the carotid artery: an evaluation with MRI.
Int J Cardiovasc Imaging 2014; 30: 91-98
- 70. Goel A, McColl R, King KS, Whittemore A, Peshock RM**
Fully automated tool to identify the aorta and compute flow using phase-contrast MRI: validation and application in a large population based study.
J Magn Reson Imaging 2014; 40: 221-228.
- 71. Iwamoto Y, Inage A, Tomlinson G, Lee KJ, Grosse-Wortmann L, Seed M, Wan A, Yoo S-J**
Direct measurement of aortic regurgitation with phase-contrast magnetic resonance is inaccurate: proposal of an alternative method of quantitation.
Pediatr Radiol 2014; 44: 1358-1369
- 72. Ait-Ali L, Siciliano V, Passino C, Molinaro S, Pasanisi E, Sicari R, Pingitore A, Festa P**
Role of stress echocardiography in operated Fallot: feasibility and detection of right ventricular response.
J Am Soc Echocardiogr 2014; 27: 1319-1328
- 73. Dragulescu A, Friedberg MK, Grosse-Wortmann L, Redington A, Mertens L**
Effect of chronic right ventricular volume overload on ventricular interaction in patients after Tetralogy of Fallot repair
J Am Soc Echocardiography 2014; 27(8): 896-902
- 74. Kowallick JT, Joseph AA, Unterberg-Buchwald C, Fasshauer M, Wijk K van, Merboldt KD, Frahm J, Lotz J, Shons JM**
Real-time phase-contrast flow MRI of the ascending aorta and superior vena cava as a function of intrathoracic pressure (Valsalva manoeuvre).
Br J Radiol 2014 (*in press*)
- 75. Das A, Wansapura JP, Gottliebson WM, Banerjee RK**
Methodology for implementing patient-specific spatial boundary condition during a cardiac cycle from phase-contrast MRI for hemodynamic assessment.
Medical Image Analysis 2014; *on-line 20 sept 2014*
- 76. Kilgour AHM, Semple S, Marshall I, Andrews P, Andrew R, Walker BR**
11 β -Hydroxysteroid dehydrogenase activity in brain does not contribute to systemic interconversion of cortisol and cortisone in healthy men

J Clin Endocrinol Metab 2014; doi: 10.1210/jc.2014-3277

- 77. Park E-A, Lee W, Kim H-K, Chung JW**
Effect of papillary muscles and trabeculae on left ventricular measurement using cardiovascular magnetic resonance imaging in patients with hypertrophic cardiomyopathy.
Korean J Radiol 2015; 16(1): doi.org/10.3348/kjr.2015.16.1.4
- 78. Hösch O, Ngyen T-T, Lauerer P, Schuster A, Kutty S, Staab S, Unterberg-Buchwald C, Sohns JM, Paul T, Lotz J, Steinmetz M**
BNP and haematological parameters are markers of severity of Ebstein's anomaly: correlation with CMR and cardiopulmonary exercise testing
Eur Heart J Cardiovasc Imaging 2015; doi/10.1093/ehjci/jeu312
- 79. Tajima Y, Suzuki E, Saito J, Murase H, Horikawa Y, Takeda J**
Elevated plasma B-type natriuretic peptide concentration and resistive index, but not decreased aortic distensibility, associate with impaired blood flow at popliteal artery in type 2 diabetic patients
Endocrine journal 2015: doi:10.1507/endocrj.EJ14-0608
- 80. Tsai-Goodman B, Zhu MY, Al-Rujaib M, Seed M, Macgowan CK**
Foetal blood flow measured using phase-contrast cardiovascular magnetic resonance – preliminary data comparing 1.5T with 3.0 T
J Magn Reson 2015: doi: 10.1186/s12968-015-0132-2
- 81. Gorter TM, Melle JP van, Freling HG, Ebels T, Bartelds B, Pieper PG, Berger RMF, Veldhuisen DJ van, Willems TP**
Pulmonary regurgitant volume is superior to fraction using background-corrected phase contrast MRI in determining the severity of regurgitation in repaired tetralogy of Fallot
Int J Cardiovasc Imaging 2015: doi: 10.1007/s10554-015-0670-6
- 82. Kröner ESJ, Grond J van der, Westenberg JJM, Wall EE van der, Siebelink H-MJ, Lamb HJ**
Morphological and functional carotid vessel wall properties in relation to cerebral white matter lesions in myocardial infarction patients.
Neth Heart J 2015; 23: 314-320.
- 83. Maroules C, Khera A, Ayers C, Goel A, Peshok RM, Abbara S, King KS**
Cardiovascular outcome associations among cardiovascular magnetic resonance measures of arterial stiffness: the Dallas heart study
J Cardiovasc Magn Reson 2014; 16: 33-42